

Herbicide Site of Action Key for Crop Injury Symptoms

To help you determine which herbicide(s) may be responsible for suspected injury on crops, this key uses the herbicide's site of action (SoA) and respective Weed Science Society of America group number; herbicides within the same SoA can cause similar symptoms. After reaching a specific SoA, you can check if any of the herbicides from that group are the source of crop response. Herbicide control (selectivity) is specified for broadleaf and/or grass weed species and remember to also observe weeds for injury symptoms.

This key is based on two traits of injury symptoms that can be used to distinguish different herbicide SoA:


Injured at emergence. The plant absorbs the herbicide from the soil either as soil-applied herbicide or carryover. The roots are normal with damaged shoots or the roots are damaged resulting in stunted plants.

Injured after emergence. The herbicide has contact activity with the older leaves, resulting in injury, or the herbicide translocates (systemic activity) to the growing points (root tips or meristems) with new tissue showing injury.

INJURED AT EMERGENCE: SOIL-APPLIED HERBICIDE OR CARRYOVER

ROOTS NORMAL, DAMAGED SHOOTS

Chlorotic / necrotic leaf margins



Broadleaves most sensitive

OR **Photosynthesis II Inhibitor (5&7)**
Atrazine, Diuron, Metribuzin, etc.

Burning / interveinal chlorosis



Burned cotyledon / hypocotyl




Broadleaves more sensitive than grasses
PPO Inhibitor (14)
Authority, Flexstar, Valor, etc.

Deformed leaves

Leafing out / buggy whipping



Drawstring



Grasses more sensitive than broadleaves
Long-chain Fatty Acid Inhibitor (15)
Dual, Outlook, Zidua, etc.

Malformed new leaves



Grasses and/or broadleaves affected
HPPD Inhibitor (27)
Balance, Callisto, Laudis, etc.

White, bleached leaves



Grasses and/or broadleaves affected
HPPD Inhibitor (27), Balance, Callisto, Laudis, etc.
Diterpene Synthesis Inhibitor (13), Command, etc.
Note that a low rate of glyphosate can also cause white, bleached leaves on seedling corn.

Leaf cupping



Broadleaves more sensitive than grasses
Synthetic Auxin (4)
2,4-D, Dicamba, Status, Stinger, etc.

ROOTS DAMAGED, STUNTED PLANTS

Clubbed root tips



Grasses most sensitive
Microtubule Inhibitor (3)
Prowl H2O, Treflan, etc.

Root proliferation
Short, thickened roots



Broadleaves more sensitive than grasses
Synthetic Auxin (4)
2,4-D, Dicamba, Status, Stinger, etc.

Pruned roots
Short, slender lateral roots



Grasses and/or broadleaves affected
ALS Inhibitor (2)
Accent, Classic, FirstRate, Pursuit, Resolve, etc.



Note that the nematicide/fungicide Fluopyran (ILevo) can mimic PPO injury symptoms (halo effect) on soybean cotyledons.

INJURED AFTER EMERGENCE: POSTEMERGENCE APPLICATION, TANK CONTAMINATION, DRIFT

CONTACT ACTIVITY: Older leaves injured, new leaves not injured

Broadleaves more sensitive than grasses



Photosynthesis II Inhibitor (5,6 & 7)
Atrazine, Basagran, Buctril, etc.



PPO Inhibitor (14)
Aim, Cobra, Flexstar, etc.

Nonselective



Glutamine Synthesis Inhibitor (10)
Glufosinate, Liberty, etc.



Photosystem I Electron Diverter (22)
Diquat, Gramoxone, etc.

TRANSLOCATING HERBICIDE WITH SYSTEMIC ACTIVITY: New leaves (meristem) injured, older leaves not injured

Leaf cupping, epinasty, leaning



New leaves chlorotic/reddish, plants stunted

Chlorotic, crinkled leaves, shortened internodes



Variable injury, chlorosis, purpling, necrosis



White, bleached leaves



Grasses and/or broadleaves affected
HPPD Inhibitor (27)
Callisto, Impact, Laudis, etc.

Chlorosis, reddened veins



Nonselective
EPSP Synthase Inhibitor (9)
Glyphosate, Roundup, etc.

Grass meristems rot



Only grasses affected
ACCase Inhibitor (1)
Assure, Poast, Select, etc.

Leaf cupping, epinasty, leaning



Broadleaves more sensitive than grasses
Synthetic Auxin (4)
2,4-D, Dicamba, Status, Stinger, etc.



Grasses and/or broadleaves affected
ALS Inhibitor (2)
Accent, Classic, Harmony, Resolve, Pursuit, etc.



Authors: Dr. Rodrigo Werle and Dr. Maxwell Coura Oliveira, Department of Agronomy, University of Wisconsin-Madison and University of Wisconsin-Extension. Revised from "Herbicide Mode of Action Key for Injury Systems" (2005) by Chris Boerboom.

An equal opportunity action employer, University of Wisconsin-Extension provides equal opportunities in employment and Programming, including Title IX requirements.

This publication is available from The Nutrient and Pest Management Program: email us at npm@hort.wisc.edu, phone: 608-265-2660, or visit our website at ipcm.wisc.edu

